

Representative Species Model: Northern Waterthrush (*Parkesia noveboracensis*)

Northern Waterthrush

Northern Waterthrush was selected as a representative species for the Designing Sustainable Landscapes project of the North Atlantic LCC (https://scholarworks.umass.edu/designing_sustainable_landscapes/). The habitat clusters (ecological systems) and associated wildlife species that it represents generally comprise of northern forest wetlands. The *Landscape Capability (LC)* index integrates habitat capability and climate suitability into a single index that reflects the relative capacity of a site to support the species.

Habitat capability (HC) - The *HC* index considers four factors representing: (1) ecological system, identifying northern forest wetlands, representing the capability of a system to provide the required invertebrate food and nesting locations needed for breeding, (2) topographic wetness, intended to identify wetland systems that have adequate hydrology conditions, (3) landscape-scale forest extent, representing the amount of undisturbed forest habitat in the landscape surrounding the homerange, and (4) large extent development, representing the effects of human-mediated landscape change that accumulate over a larger geographical area and that may penetrate more deeply into the forest than the processes of local edge effects, such as population increases of cowbirds and generalist predators. The *HC* index represents the relative capacity of a site to provide the habitat needed by the species based on current scientific knowledge.


Climate niche (CN) - The *CN* index uses two climate variables representing: (1) growing degree days, (2) annual precipitation, (3) growing season precipitation, (4) maximum summer temperature, (5) minimum winter temperature, and (6) annual temperature and is based on a statistical model derived from 5,522




Breeding Bird Survey (BBS) route segments distributed through the Humid Temperate Domain. The *CN* index represents the probability of the climate being suitable for the species based on its current distribution in relation to current climate.

Landscape Capability (LC) - The *LC* index is computed as the product of the *HC* and *CN* indices (see map). Thus, the index computed for 2010 reflects the gradient of worst (0) to best (maximum value) sites within the landscape that support this species during the breeding season. Note, we also compute this index for the future (e.g., 2080) based on output from the landscape change model.

Model performance was evaluated using an independent dataset (eBird occurrence data; 1,243 present locations and 2,517 absent locations) and determined to be acceptable (Kappa = 0.81, Deviance explained=68%, AUC = 0.93).





See technical document on species at https://scholarworks.umass.edu/designing_sustainable_landscapes/ for a detailed description of the Landscape Capability modeling process.